



Self-adjoint extensions of differential operators on Riemannian manifolds

Ognjen Milatovic, Francoise Truc

► To cite this version:

Ognjen Milatovic, Francoise Truc. Self-adjoint extensions of differential operators on Riemannian manifolds. *Annals of Global Analysis and Geometry*, In press, 49 (1), pp.87-103. 10.1007/s10455-015-9482-0 . hal-01149461

HAL Id: hal-01149461

<https://hal.science/hal-01149461>

Submitted on 7 May 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SELF-ADJOINT EXTENSIONS OF DIFFERENTIAL OPERATORS ON RIEMANNIAN MANIFOLDS

OGNJEN MILATOVIC, FRANÇOISE TRUC

ABSTRACT. We study $H = D^*D + V$, where D is a first order elliptic differential operator acting on sections of a Hermitian vector bundle over a Riemannian manifold M , and V is a Hermitian bundle endomorphism. In the case when M is geodesically complete, we establish the essential self-adjointness of positive integer powers of H . In the case when M is not necessarily geodesically complete, we give a sufficient condition for the essential self-adjointness of H , expressed in terms of the behavior of V relative to the Cauchy boundary of M .

1. INTRODUCTION

As a fundamental problem in mathematical physics, self-adjointness of Schrödinger operators has attracted the attention of researchers over many years now, resulting in numerous sufficient conditions for this property in $L^2(\mathbb{R}^n)$. For reviews of the corresponding results, see, for instance, the books [14, 29].

The study of the corresponding problem in the context of a non-compact Riemannian manifold was initiated by Gaffney [15, 16] with the proof of the essential self-adjointness of the Laplacian on differential forms. About two decades later, Cordes (see Theorem 3 in [11]) proved the essential self-adjointness of positive integer powers of the operator

$$\Delta_{M,\mu} := -\frac{1}{\kappa} \left(\frac{\partial}{\partial x^i} \left(\kappa g^{ij} \frac{\partial}{\partial x^j} \right) \right) \quad (1.1)$$

on an n -dimensional geodesically complete Riemannian manifold M equipped with a (smooth) metric $g = (g_{ij})$ (here, $(g^{ij}) = ((g_{ij})^{-1})$) and a positive smooth measure $d\mu$ (i.e. in any local coordinates x^1, x^2, \dots, x^n there exists a strictly positive C^∞ -density $\kappa(x)$ such that $d\mu = \kappa(x) dx^1 dx^2 \dots dx^n$). Theorem 1 of our paper extends this result to the operator $(D^*D + V)^k$ for all $k \in \mathbb{Z}_+$, where D is a first order elliptic differential operator acting on sections of a Hermitian vector bundle over a geodesically complete Riemannian manifold, D^* is the formal adjoint of D , and V is a self-adjoint Hermitian bundle endomorphism; see Section 2.3 for details.

In the context of a general Riemannian manifold (not necessarily geodesically complete), Cordes (see Theorem IV.1.1 in [12] and Theorem 4 in [11]) proved the essential self-adjointness of P^k for all $k \in \mathbb{Z}_+$, where

$$Pu := \Delta_{M,\mu}u + qu, \quad u \in C^\infty(M), \quad (1.2)$$

and $q \in C^\infty(M)$ is real-valued. Thanks to a Roelcke-type estimate (see Lemma 3.1 below), the technique of Cordes [12] can be applied to the operator $(D^*D + V)^k$ acting on sections of Hermitian vector bundles over a general Riemannian manifold. To make our exposition shorter, in Theorem 1 we consider the geodesically complete case. Our Theorem 2 concerns $(\nabla^*\nabla + V)^k$, where ∇ is a metric connection on a Hermitian vector bundle over a non-compact geodesically complete Riemannian manifold. This result extends Theorem 1.1 of [13] where Cordes showed that if (M, g) is non-compact and geodesically complete and P is semi-bounded from below on $C_c^\infty(M)$, then P^k is essentially self-adjoint on $C_c^\infty(M)$, for all $k \in \mathbb{Z}_+$.

For the remainder of the introduction, the notation $D^*D + V$ is used in the same sense as described earlier in this section. In the setting of geodesically complete Riemannian manifolds, the essential self-adjointness of $D^*D + V$ with $V \in L_{\text{loc}}^\infty$ was established in [21], providing a generalization of the results in [3, 27, 28, 32] concerning Schrödinger operators on functions (or differential forms). Subsequently, the operator $D^*D + V$ with a singular potential V was considered in [5]. Recently, in the case $V \in L_{\text{loc}}^\infty$, the authors of [4] extended the main result of [5] to the operator $D^*D + V$ acting on sections of infinite-dimensional bundles whose fibers are modules of finite type over a von Neumann algebra.

In the context of an incomplete Riemannian manifold, the authors of [17, 22, 23] studied the so-called Gaffney Laplacian, a self-adjoint realization of the scalar Laplacian generally different from the closure of $\Delta_{M, d\mu}|_{C_c^\infty(M)}$. For a study of Gaffney Laplacian on differential forms, see [24].

Our Theorem 3 gives a condition on the behavior of V relative to the Cauchy boundary of M that will guarantee the essential self-adjointness of $D^*D + V$; for details see Section 2.4 below. Related results can be found in [6, 25, 26] in the context of (magnetic) Schrödinger operators on domains in \mathbb{R}^n , and in [10] concerning the magnetic Laplacian on domains in \mathbb{R}^n and certain types of Riemannian manifolds.

Finally, let us mention that Chernoff [7] used the hyperbolic equation approach to establish the essential self-adjointness of positive integer powers of Laplace–Beltrami operator on differential forms. This approach was also applied in [2, 8, 9, 18, 19, 31] to prove essential self-adjointness of second-order operators (acting on scalar functions or sections of Hermitian vector bundles) on Riemannian manifolds. Additionally, the authors of [18, 19] used path integral techniques.

The paper is organized as follows. The main results are stated in Section 2, a preliminary lemma is proven in Section 3, and the main results are proven in Sections 4–6.

2. MAIN RESULTS

2.1. The setting. Let M be an n -dimensional smooth, connected Riemannian manifold without boundary. We denote the Riemannian metric on M by g^{TM} . We assume that M is equipped with a positive smooth measure $d\mu$, i.e. in any local coordinates x^1, x^2, \dots, x^n there exists a strictly positive C^∞ -density $\kappa(x)$ such that $d\mu = \kappa(x) dx^1 dx^2 \dots dx^n$. Let E be a Hermitian vector bundle over M and let $L^2(E)$ denote the Hilbert space of square integrable sections of E with respect to the inner product

$$(u, v) = \int_M \langle u(x), v(x) \rangle_{E_x} d\mu(x), \quad (2.1)$$

where $\langle \cdot, \cdot \rangle_{E_x}$ is the fiberwise inner product. The corresponding norm in $L^2(E)$ is denoted by $\| \cdot \|$. In Sobolev space notations $W_{\text{loc}}^{k,2}(E)$ used in this paper, the superscript $k \in \mathbb{Z}_+$ indicates the order of the highest derivative. The corresponding dual space is denoted by $W_{\text{loc}}^{-k,2}(E)$.

Let F be another Hermitian vector bundle on M . We consider a first order differential operator $D: C_c^\infty(E) \rightarrow C_c^\infty(F)$, where C_c^∞ stands for the space of smooth compactly supported sections. In the sequel, by $\sigma(D)$ we denote the principal symbol of D .

Assumption (A0) Assume that D is elliptic. Additionally, assume that there exists a constant $\lambda_0 > 0$ such that

$$|\sigma(D)(x, \xi)| \leq \lambda_0 |\xi|, \quad \text{for all } x \in M, \xi \in T_x^*M, \quad (2.2)$$

where $|\xi|$ is the length of ξ induced by the metric g^{TM} and $|\sigma(D)(x, \xi)|$ is the operator norm of $\sigma(D)(x, \xi): E_x \rightarrow F_x$.

Remark 2.2. Assumption (A0) is satisfied if $D = \nabla$, where $\nabla: C^\infty(E) \rightarrow C^\infty(T^*M \otimes E)$ is a covariant derivative corresponding to a metric connection on a Hermitian vector bundle E over M .

2.3. Schrödinger-type Operator. Let $D^*: C_c^\infty(F) \rightarrow C_c^\infty(E)$ be the formal adjoint of D with respect to the inner product (2.1). We consider the operator

$$H = D^*D + V, \quad (2.3)$$

where $V \in L_{\text{loc}}^\infty(\text{End } E)$ is a linear self-adjoint bundle endomorphism. In other words, for all $x \in M$, the operator $V(x): E_x \rightarrow E_x$ is self-adjoint and $|V(x)| \in L_{\text{loc}}^\infty(M)$, where $|V(x)|$ is the norm of the operator $V(x): E_x \rightarrow E_x$.

2.4. Statements of Results.

Theorem 1. *Let M , g^{TM} , and $d\mu$ be as in Section 2.1. Assume that (M, g^{TM}) is geodesically complete. Let E and F be Hermitian vector bundles over M , and let $D: C_c^\infty(E) \rightarrow C_c^\infty(F)$ be a first order differential operator satisfying the assumption (A0). Assume that $V \in C^\infty(\text{End } E)$ and*

$$V(x) \geq C, \quad \text{for all } x \in M,$$

where C is a constant, and the inequality is understood in operator sense. Then H^k is essentially self-adjoint on $C_c^\infty(E)$, for all $k \in \mathbb{Z}_+$.

Remark 2.5. In the case $V = 0$, the following result related to Theorem 1 can be deduced from Chernoff (see Theorem 2.2 in [7]):

Assume that (M, g) is a geodesically complete Riemannian manifold with metric g . Let D be as in Theorem 1, and define

$$c(x) := \sup\{|\sigma(D)(x, \xi)|: |\xi|_{T_x^*M} = 1\}.$$

Fix $x_0 \in M$ and define

$$c(r) := \sup_{x \in B(x_0, r)} c(x),$$

where $r > 0$ and $B(x_0, r) := \{x \in M : d_g(x_0, x) < r\}$. Assume that

$$\int_0^\infty \frac{1}{c(r)} dr = \infty. \quad (2.4)$$

Then the operator $(D^*D)^k$ is essentially self-adjoint on $C_c^\infty(E)$ for all $k \in \mathbb{Z}_+$.

At the end of this section we give an example of an operator for which Theorem 1 guarantees the essential self-adjointness of $(D^*D)^k$, whereas Chernoff's result cannot be applied.

The next theorem is concerned with operators whose potential V is not necessarily semi-bounded from below.

Theorem 2. Let M , g^{TM} , and $d\mu$ be as in Section 2.1. Assume that (M, g^{TM}) is noncompact and geodesically complete. Let E be a Hermitian vector bundle over M and let ∇ be a Hermitian connection on E . Assume that $V \in C^\infty(\text{End } E)$ and

$$V(x) \geq q(x), \quad \text{for all } x \in M, \quad (2.5)$$

where $q \in C^\infty(M)$ and the inequality is understood in the sense of operators $E_x \rightarrow E_x$. Additionally, assume that

$$((\Delta_{M,\mu} + q)u, u) \geq C\|u\|^2, \quad \text{for all } u \in C_c^\infty(M), \quad (2.6)$$

where $C \in \mathbb{R}$ and $\Delta_{M,\mu}$ is as in (1.1) with g replaced by g^{TM} . Then the operator $(\nabla^*\nabla + V)^k$ is essentially self-adjoint on $C_c^\infty(E)$, for all $k \in \mathbb{Z}_+$.

Remark 2.6. Let us stress that non-compactness is required in the proof to ensure the existence of a positive smooth solution of an equation involving $\Delta_{M,\mu} + q$. In the case of a compact manifold, such a solution exists under an additional assumption; see Theorem III.6.3 in [12].

In our last result we will need the notion of Cauchy boundary. Let $d_{g^{TM}}$ be the distance function corresponding to the metric g^{TM} . Let $(\widehat{M}, \widehat{d}_{g^{TM}})$ be the metric completion of $(M, d_{g^{TM}})$. We define the *Cauchy boundary* $\partial_C M$ as follows: $\partial_C M := \widehat{M} \setminus M$. Note that $(M, d_{g^{TM}})$ is metrically complete if and only if $\partial_C M$ is empty. For $x \in M$ we define

$$r(x) := \inf_{z \in \partial_C M} \widehat{d}_{g^{TM}}(x, z). \quad (2.7)$$

We will also need the following assumption:

Assumption (A1) Assume that \widehat{M} is a smooth manifold and that the metric g^{TM} extends to $\partial_C M$.

Remark 2.7. Let N be a (smooth) n -dimensional Riemannian manifold without boundary. Denote the metric on N by g^{TN} and assume that (N, g^{TN}) is geodesically complete. Let Σ be a k -dimensional closed sub-manifold of N with $k < n$. Then $M := N \setminus \Sigma$ has the properties $\widehat{M} = N$ and $\partial_C M = \Sigma$. Thus, assumption (A1) is satisfied.

Theorem 3. Let M , g^{TM} , and $d\mu$ be as in Section 2.1. Assume that (A1) is satisfied. Let E and F be Hermitian vector bundles over M , and let $D: C_c^\infty(E) \rightarrow C_c^\infty(F)$ be a first order differential operator satisfying the assumption (A0). Assume that $V \in L_{\text{loc}}^\infty(\text{End } E)$ and there exists a constant C such that

$$V(x) \geq \left(\frac{\lambda_0}{r(x)} \right)^2 - C, \quad \text{for all } x \in M, \quad (2.8)$$

where λ_0 is as in (2.2), the distance $r(x)$ is as in (2.7), and the inequality is understood in the sense of linear operators $E_x \rightarrow E_x$. Then H is essentially self-adjoint on $C_c^\infty(E)$.

In order to describe the example mentioned in Remark 2.5, we need the following

Remark 2.8. As explained in [5], we can use a first-order elliptic operator $D: C_c^\infty(E) \rightarrow C_c^\infty(F)$ to define a metric on M . For $\xi, \eta \in T_x^*M$, define

$$\langle \xi, \eta \rangle = \frac{1}{m} \text{Re Tr} ((\sigma(D)(x, \xi))^* \sigma(D)(x, \eta)), \quad m = \dim E_x, \quad (2.9)$$

where Tr denotes the usual trace of a linear operator. Since D is an elliptic first-order differential operator and $\sigma(D)(x, \xi)$ is linear in ξ , it is easily checked that (2.9) defines an inner product on T_x^*M . Its dual defines a Riemannian metric on M . Denoting this metric by g^{TM} and using elementary linear algebra, it follows that (2.2) is satisfied with $\lambda_0 = \sqrt{m}$.

Example 2.9. Let $M = \mathbb{R}^2$ with the standard metric and measure, and $V = 0$. Denoting respectively by $C_c^\infty(\mathbb{R}^2; \mathbb{R})$ and $C_c^\infty(\mathbb{R}^2; \mathbb{R}^2)$ the spaces of smooth compactly supported functions $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ and $f: \mathbb{R}^2 \rightarrow \mathbb{R}^2$, we define the operator $D: C_c^\infty(\mathbb{R}^2; \mathbb{R}) \rightarrow C_c^\infty(\mathbb{R}^2; \mathbb{R}^2)$ by

$$D = \begin{pmatrix} a(x, y) \frac{\partial}{\partial x} \\ b(x, y) \frac{\partial}{\partial y} \end{pmatrix},$$

where

$$\begin{aligned} a(x, y) &= (1 - \cos(2\pi e^x))x^2 + 1; \\ b(x, y) &= (1 - \sin(2\pi e^y))y^2 + 1. \end{aligned}$$

Since a, b are smooth real-valued nowhere vanishing functions in \mathbb{R}^2 , it follows that the operator D is elliptic. We are interested in the operator

$$H := D^*D = -\frac{\partial}{\partial x} \left(a^2 \frac{\partial}{\partial x} \right) - \frac{\partial}{\partial y} \left(b^2 \frac{\partial}{\partial y} \right).$$

The matrix of the inner product on T^*M defined by D via (2.9) is $\text{diag}(a^2/2, b^2/2)$. The matrix of the corresponding Riemannian metric g^{TM} on M is $\text{diag}(2a^{-2}, 2b^{-2})$, so the metric itself is $ds^2 = 2a^{-2}dx^2 + 2b^{-2}dy^2$ and it is geodesically complete (see Example 3.1 of [5]). Moreover, thanks to Remark 2.8, assumption (A0) is satisfied. Thus, by Theorem 1 the operator $(D^*D)^k$ is essentially self-adjoint for all $k \in \mathbb{Z}_+$. Furthermore, in Example 3.1 of [5] it was shown that for the considered operator D the condition (2.4) is not satisfied. Thus, the result stated in Remark 2.5 does not apply.

3. ROELCKE-TYPE INEQUALITY

Let M , $d\mu$, D , and $\sigma(D)$ be as in Section 2.1. Set $\widehat{D} := -i\sigma(D)$, where $i = \sqrt{-1}$. Then for any Lipschitz function $\psi: M \rightarrow \mathbb{R}$ and $u \in W_{\text{loc}}^{1,2}(E)$ we have

$$D(\psi u) = \widehat{D}(d\psi)u + \psi Du, \quad (3.1)$$

where we have suppressed x for simplicity. We also note that $\widehat{D}^*(\xi) = -(\widehat{D}(\xi))^*$, for all $\xi \in T_x^*M$.

For a compact set $K \subset M$, and $u, v \in W_{\text{loc}}^{1,2}(E)$, we define

$$(u, v)_K := \int_K \langle u(x), v(x) \rangle d\mu(x), \quad (Du, Dv)_K := \int_K \langle Du(x), Dv(x) \rangle d\mu(x). \quad (3.2)$$

In order to prove Theorem 1 we need the following important lemma, which is an extension of Lemma 2.1 in [12] to operator (2.3). In the context of the scalar Laplacian on a Riemannian manifold, this kind of result is originally due to Roelcke [30].

Lemma 3.1. *Let M , g^{TM} , and $d\mu$ be as in Section 2.1. Let E and F be Hermitian vector bundles over M , and let $D: C_c^\infty(E) \rightarrow C_c^\infty(F)$ be a first order differential operator satisfying the assumption (A0). Let $\rho: M \rightarrow [0, \infty)$ be a function satisfying the following properties:*

- (i) $\rho(x)$ is Lipschitz continuous with respect to the distance induced by the metric g^{TM} ;
- (ii) $\rho(x_0) = 0$, for some fixed $x_0 \in M$;
- (iii) the set $B_T := \{x \in M: \rho(x) \leq T\}$ is compact, for some $T > 0$.

Then the following inequality holds for all $u \in W_{\text{loc}}^{2,2}(E)$ and $v \in W_{\text{loc}}^{2,2}(E)$:

$$\int_0^T |(Du, Dv)_{B_t} - (D^*Du, v)_{B_t}| dt \leq \lambda_0 \int_{B_T} |d\rho(x)| |Du(x)| |v(x)| d\mu(x), \quad (3.3)$$

where B_t is as in (iii) (with t instead of T), the constant λ_0 is as in (2.2), and $|d\rho(x)|$ is the length of $d\rho(x) \in T_x^*M$ induced by g^{TM} .

Proof. For $\varepsilon > 0$ and $t \in (0, T)$, we define a continuous piecewise linear function $F_{\varepsilon,t}$ as follows:

$$F_{\varepsilon,t}(s) = \begin{cases} 1 & \text{for } s < t - \varepsilon \\ (t - s)/\varepsilon & \text{for } t - \varepsilon \leq s < t \\ 0 & \text{for } s \geq t \end{cases}$$

The function $f_{\varepsilon,t}(x) := F_{\varepsilon,t}(\rho(x))$, is Lipschitz continuous with respect to the distance induced by the metric g^{TM} , and $d(f_{\varepsilon,t}(\rho(x))) = (F'_{\varepsilon,t}(\rho(x)))d\rho(x)$. Moreover we have $f_{\varepsilon,t}v \in W_{\text{loc}}^{1,2}(E)$ for all $v \in W_{\text{loc}}^{1,2}(E)$, since

$$D(f_{\varepsilon,t}v) = \widehat{D}(df_{\varepsilon,t})v + f_{\varepsilon,t}Dv.$$

It follows from the compactness of B_T that B_t is compact for all $t \in (0, T)$. Using integration by parts (see Lemma 8.8 in [5]), for all $u \in W_{\text{loc}}^{2,2}(E)$ and $v \in W_{\text{loc}}^{2,2}(E)$ we have

$$(D^*Du, v f_{\varepsilon,t})_{B_t} = (Du, D(v f_{\varepsilon,t}))_{B_t} = (Du, f_{\varepsilon,t}Dv)_{B_t} + (Du, \widehat{D}(df_{\varepsilon,t})v)_{B_t},$$

which, together with (2.2), gives

$$\begin{aligned}
& |(Du, f_{\varepsilon,t}Dv)_{B_t} - (D^*Du, v f_{\varepsilon,t})_{B_t}| = |(Du, \widehat{D}(df_{\varepsilon,t})v)_{B_t}| \\
& \leq \int_{B_t} |Du(x)| |\widehat{D}(df_{\varepsilon,t}(x))v(x)| d\mu(x) \leq \lambda_0 \int_{B_t} |Du(x)| |df_{\varepsilon,t}(x)| |v(x)| d\mu(x) \\
& = \lambda_0 \int_{B_t} |Du(x)| |F'_{\varepsilon,t}(\rho(x))| |d\rho(x)| |v(x)| d\mu(x) \\
& \leq \lambda_0 \int_{B_T} |Du(x)| |F'_{\varepsilon,t}(\rho(x))| |d\rho(x)| |v(x)| d\mu(x), \tag{3.4}
\end{aligned}$$

where $|df_{\varepsilon,t}(x)|$ and $|d\rho(x)|$ are the norms of $df_{\varepsilon,t}(x) \in T_x^*M$ and $d\rho(x) \in T_x^*M$ induced by g^{TM} .

Fixing $\varepsilon > 0$, integrating the leftmost and the rightmost side of (3.4) from $t = 0$ to $t = T$, and noting that $F'_{\varepsilon,t}(\rho(x))$ is the only term on the rightmost side depending on t , we obtain

$$\begin{aligned}
& \int_0^T |(Du, f_{\varepsilon,t}Dv)_{B_t} - (D^*Du, v f_{\varepsilon,t})_{B_t}| dt \\
& \leq \lambda_0 \int_{B_T} |Du(x)| |d\rho(x)| |v(x)| I_\varepsilon(x) d\mu(x), \tag{3.5}
\end{aligned}$$

where

$$I_\varepsilon(x) := \int_0^T |F'_{\varepsilon,t}(\rho(x))| dt.$$

We now let $\varepsilon \rightarrow 0+$ in (3.5). On the left-hand side of (3.5), as $\varepsilon \rightarrow 0+$, we have $f_{\varepsilon,t}(x) \rightarrow \chi_{B_t}(x)$ almost everywhere, where $\chi_{B_t}(x)$ is the characteristic function of the set B_t . Additionally, $|f_{\varepsilon,t}(x)| \leq 1$ for all $x \in B_t$ and all $t \in (0, T)$; thus, by dominated convergence theorem, as $\varepsilon \rightarrow 0+$ the left-hand side of (3.5) converges to the left-hand side of (3.3). On the right-hand side of (3.5) an easy calculation shows that $I_\varepsilon(x) \rightarrow 1$, as $\varepsilon \rightarrow 0+$. Additionally, we have $|I_\varepsilon(x)| \leq 1$, a.e. on B_T ; hence, by the dominated convergence theorem, as $\varepsilon \rightarrow 0+$ the right-hand side of (3.5) converges to the right-hand side of (3.3). This establishes the inequality (3.3). \square

4. PROOF OF THEOREM 1

We first give the definitions of minimal and maximal operators associated with the expression H in (2.3).

4.1. Minimal and Maximal Operators. We define $H_{\min}u := Hu$, with $\text{Dom}(H_{\min}) := C_c^\infty(E)$, and $H_{\max} := (H_{\min})^*$, where T^* denotes the adjoint of operator T . Denoting $\mathcal{D}_{\max} := \{u \in L^2(E) : Hu \in L^2(E)\}$, we recall the following well-known property: $\text{Dom}(H_{\max}) = \mathcal{D}_{\max}$ and $H_{\max}u = Hu$ for all $u \in \mathcal{D}_{\max}$.

From now on, throughout this section, we assume that the hypotheses of Theorem 1 are satisfied. Let $x_0 \in M$, and define $\rho(x) := d_{g^{TM}}(x_0, x)$, where $d_{g^{TM}}$ is the distance function corresponding to the metric g^{TM} . By the definition of $\rho(x)$ and the geodesic completeness of (M, g^{TM}) , it follows that $\rho(x)$ satisfies all hypotheses of Lemma 3.1. Using Lemma 3.1 and Proposition 4.2 below, we are able to apply the method of Cordes [11, 12] to our context. As we

will see, Cordes's technique reduces our problem to a system of ordinary differential inequalities of the same type as in Section IV.3 of [12].

Proposition 4.2. *Let A be a densely defined operator with domain \mathcal{D} in a Hilbert space \mathcal{H} . Assume that A is semi-bounded from below, that $A\mathcal{D} \subseteq \mathcal{D}$, and that there exists $c_0 \in \mathbb{R}$ such that the following two properties hold:*

- (i) $((A + c_0 I)u, u)_{\mathcal{H}} \geq \|u\|_{\mathcal{H}}^2$, for all $u \in \mathcal{D}$, where I denotes the identity operator in \mathcal{H} ;
- (ii) $(A + c_0 I)^k$ is essentially self-adjoint on \mathcal{D} , for some $k \in \mathbb{Z}_+$.

Then, $(A + cI)^j$ is essentially self-adjoint on \mathcal{D} , for all $j = 1, 2, \dots, k$ and all $c \in \mathbb{R}$.

Remark 4.3. To prove Proposition 4.2, one may mimick the proof of Proposition 1.4 in [12], which was carried out for the operator P defined in (1.2) with $\mathcal{D} = C_c^\infty(M)$, since only abstract functional analysis facts and the property $P\mathcal{D} \subseteq \mathcal{D}$ were used.

We start the proof of Theorem 1 by noticing that the operator H_{\min} is essentially self-adjoint on $C_c^\infty(E)$; see Corollary 2.9 in [5]. Thanks to Proposition 4.2, without any loss of generality we can change $V(x)$ to $V(x) + C \text{Id}(x)$, where C is a sufficiently large constant in order to have

$$V(x) \geq (\lambda_0^2 + 1)\text{Id}(x), \quad \text{for all } x \in M, \quad (4.1)$$

where λ_0 is as in (2.2) and $\text{Id}(x)$ is the identity endomorphism of E_x . Using non-negativity of D^*D and (4.1) we have

$$(H_{\min}u, u) \geq \|u\|^2, \quad \text{for all } u \in C_c^\infty(E), \quad (4.2)$$

which leads to

$$\|u\|^2 \leq (Hu, u) \leq \|Hu\|\|u\|, \quad \text{for all } u \in C_c^\infty(E),$$

and, hence, $\|Hu\| \geq \|u\|$, for all $u \in C_c^\infty(E)$. Therefore,

$$(H^2u, u) = (Hu, Hu) = \|Hu\|^2 \geq \|u\|^2, \quad \text{for all } u \in C_c^\infty(E), \quad (4.3)$$

and

$$(H^3u, u) = (HHu, Hu) \geq \|Hu\|^2 \geq \|u\|^2, \quad \text{for all } u \in C_c^\infty(E).$$

By (4.3) we have

$$\|u\|^2 \leq (H^2u, u) \leq \|H^2u\|\|u\|, \quad \text{for all } u \in C_c^\infty(E),$$

and, hence, $\|H^2u\| \geq \|u\|$, for all $u \in C_c^\infty(E)$. This, in turn, leads to

$$(H^4u, u) = (H^2u, H^2u) = \|H^2u\|^2 \geq \|u\|^2, \quad \text{for all } u \in C_c^\infty(E).$$

Continuing like this, we obtain $(H^k u, u) \geq \|u\|^2$, for all $u \in C_c^\infty(E)$ and all $k \in \mathbb{Z}_+$. In this case, by an abstract fact (see Theorem X.26 in [29]), the essential self-adjointness of H^k on $C_c^\infty(E)$ is equivalent to the following statement: if $u \in L^2(E)$ satisfies $H^k u = 0$, then $u = 0$.

Let $u \in L^2(E)$ satisfy $H^k u = 0$. Since $V \in C^\infty(E)$, by local elliptic regularity it follows that $u \in C^\infty(E) \cap L^2(E)$. Define

$$f_j := H^{k-j}u, \quad j = 0, \pm 1, \pm 2, \dots \quad (4.4)$$

Here, in the case $k - j < 0$, the definition (4.4) is interpreted as $((H_{\max})^{-1})^{j-k}$. We already noted that H_{\min} is essentially self-adjoint and positive. Furthermore, it is well known that the self-adjoint closure of H_{\min} coincides with H_{\max} . Therefore H_{\max} is a positive self-adjoint operator, and $(H_{\max})^{-1}: L^2(E) \rightarrow L^2(E)$ is bounded. This, together with $f_k = u \in L^2(E)$ explains the following property: $f_j \in L^2(E)$, for all $j \geq k$. Additionally, observe that $f_j = 0$ for all $j \leq 0$ because $f_0 = 0$. Furthermore, we note that $f_j \in C^\infty(E)$, for all $j \in \mathbb{Z}$. The last assertion is obvious for $j \leq k$, and for $j > k$ it can be seen by showing that $H^j f_j = 0$ in distributional sense and using $f_j \in L^2(E)$ together with local elliptic regularity. To see this, let $v \in C_c^\infty(E)$ be arbitrary, and note that

$$(f_j, H^j v) = (H^{k-j} u, H^j v) = (u, H^k v) = (H^k u, v) = 0.$$

Finally, observe that

$$H^l f_j = f_{j-l}, \quad \text{for all } j \in \mathbb{Z} \text{ and } l \in \mathbb{Z}_+ \cup \{0\}. \quad (4.5)$$

With f_j as in (4.4), define the functions α_j and β_j on the interval $0 \leq T < \infty$ by the formulas

$$\alpha_j(T) := \lambda_0^2 \int_0^T (f_j, f_j)_{B_t} dt, \quad \beta_j(T) := \int_0^T (Df_j, Df_j)_{B_t} dt, \quad (4.6)$$

where λ_0 is as in (4.1) and $(\cdot, \cdot)_{B_t}$ is as in (3.2).

In the sequel, to simplify the notations, the functions $\alpha_j(T)$ and $\beta_j(T)$, the inner products $(\cdot, \cdot)_{B_t}$, and the corresponding norms $\|\cdot\|_{B_t}$ appearing in (4.6) will be denoted by α_j , β_j , $(\cdot, \cdot)_t$, and $\|\cdot\|_t$, respectively.

Note that α_j and β_j are absolutely continuous on $[0, \infty)$. Furthermore, α_j and β_j have a left first derivative and a right first derivative at each point. Additionally, α_j and β_j are differentiable, except at (at most) countably many points. In the sequel, to simplify notations, we shall denote the right first derivatives of α_j and β_j by α'_j and β'_j . Note that α_j , β_j , α'_j and β'_j are non-decreasing and non-negative functions. Note also that α_j and β_j are convex functions. Furthermore, since $f_j = 0$ for all $j \leq 0$, it follows that $\alpha_j \equiv 0$ and $\beta_j \equiv 0$ for all $j \leq 0$. Finally, using (4.1) and the property $f_j \in L^2(E) \cap C^\infty(E)$ for all $j \geq k$, observe that

$$\lambda_0^2(f_j, f_j) + (Df_j, Df_j) \leq (Vf_j, f_j) + (Df_j, Df_j) = (f_j, Hf_j) = (f_j, f_{j-1}) < \infty,$$

for all $j > k$. Here, “integration by parts” in the first equality is justified because H_{\min} is essentially self-adjoint (i.e. $C_c^\infty(E)$ is an operator core of H_{\max}). Hence, α'_j and β'_j are bounded for all $j > k$. It turns out that α_j and β_j satisfy a system of differential inequalities, as seen in the next proposition.

Proposition 4.4. *Let α_j and β_j be as in (4.6). Then, for all $j \geq 1$ and all $T \geq 0$ we have*

$$\alpha_j + \beta_j \leq \sqrt{\alpha'_j \beta'_j} + \sum_{l=0}^{\infty} \left(\sqrt{\alpha'_{j+l+1} \beta'_{j-l-1}} + \sqrt{\alpha'_{j-l-1} \beta'_{j+l+1}} \right) \quad (4.7)$$

and

$$\alpha_j \leq \lambda_0^2 \left(\sum_{l=0}^{\infty} \left(\sqrt{\alpha'_{j+l+1} \beta'_{j-l-1}} + \sqrt{\alpha'_{j-l-1} \beta'_{j+l+1}} \right) \right), \quad (4.8)$$

where λ_0 is as in (4.1) and α'_i, β'_i denote the right-hand derivatives.

Remark 4.5. Note that the sums in (4.7) and (4.8) are finite since $\alpha_i \equiv 0$ and $\beta_i \equiv 0$ for $i \leq 0$. As our goal is to show that $f_k = u = 0$, we will only use the first k inequalities in (4.7) and the first k inequalities in (4.8).

Proof of Proposition 4.4. From (4.6) and (4.1) it follows that

$$\alpha_j + \beta_j \leq \int_0^T ((f_j, Vf_j)_t + (Df_j, Df_j)_t) dt. \quad (4.9)$$

We start from (4.9), use (3.3), Cauchy–Schwarz inequality, and (4.5) to obtain

$$\begin{aligned} \alpha_j + \beta_j &\leq \int_0^T ((f_j, Vf_j)_t + (Df_j, Df_j)_t) dt \\ &= \int_0^T |(f_j, Hf_j)_t - (f_j, D^*Df_j)_t + (Df_j, Df_j)_t| dt \\ &\leq \lambda_0 \int_{B_T} |Df_j(x)| |f_j(x)| d\mu(x) + \int_0^T |(f_j, Hf_j)_t| dt \\ &\leq \sqrt{\alpha'_j \beta'_j} + \int_0^T |(Hf_{j+1}, f_{j-1})_t| dt. \end{aligned}$$

We continue the process as follows:

$$\begin{aligned} \alpha_j + \beta_j &\leq \sqrt{\alpha'_j \beta'_j} + \int_0^T |(Hf_{j+1}, f_{j-1})_t| dt \\ &= \sqrt{\alpha'_j \beta'_j} + \int_0^T |(D^*Df_{j+1}, f_{j-1})_t + (f_{j+1}, Vf_{j-1})_t| dt \\ &\leq \sqrt{\alpha'_j \beta'_j} + \int_0^T |(D^*Df_{j+1}, f_{j-1})_t - (Df_{j+1}, Df_{j-1})_t| dt \\ &\quad + \int_0^T |(Df_{j+1}, Df_{j-1})_t - (f_{j+1}, D^*Df_{j-1})_t| dt + \int_0^T |(f_{j+1}, Hf_{j-1})_t| dt \\ &\leq \sqrt{\alpha'_j \beta'_j} + \sqrt{\alpha'_{j+1} \beta'_{j-1}} + \sqrt{\alpha'_{j-1} \beta'_{j+1}} + \int_0^T |(Hf_{j+2}, f_{j-2})_t| dt, \end{aligned}$$

where we used triangle inequality, (3.3), Cauchy–Schwarz inequality, and (4.5). We continue like this until the last term reaches the subscript $j - l \leq 0$, which makes the last term equal zero by properties of f_i discussed above. This establishes (4.7).

To show (4.8), we start from the definition of α_j , use (3.3), Cauchy–Schwarz inequality, and (4.5) to obtain

$$\begin{aligned}
\alpha_j &= \lambda_0^2 \int_0^T (f_j, f_j)_t dt = \lambda_0^2 \int_0^T |(f_j, H f_{j+1})_t| dt \\
&= \lambda_0^2 \int_0^T |(f_j, D^* D f_{j+1})_t + (V f_j, f_{j+1})_t| dt \\
&\leq \lambda_0^2 \int_0^T |(f_j, D^* D f_{j+1})_t - (D f_j, D f_{j+1})_t| dt \\
&\quad + \lambda_0^2 \int_0^T |(D f_j, D f_{j+1})_t - (D^* D f_j, f_{j+1})_t| dt + \lambda_0^2 \int_0^T |(H f_j, f_{j+1})_t| dt \\
&\leq \lambda_0^2 \left(\sqrt{\alpha'_{j+1} \beta'_j} + \sqrt{\alpha'_j \beta'_{j+1}} \right) + \lambda_0^2 \int_0^T |(f_{j-1}, f_{j+1})_t| dt.
\end{aligned}$$

We continue like this until the last term reaches the subscript $j - l \leq 0$, which makes the last term equal zero by properties of f_i discussed above. This establishes (4.8). \square

End of the proof of Theorem 1. We will now transform the system (4.7)–(4.8) by introducing new variables:

$$\omega_j(T) := \alpha_j(T) + \beta_j(T), \quad \theta_j(T) := \alpha_j(T) - \beta_j(T) \quad T \in [0, \infty). \quad (4.10)$$

To carry out the transformation, observe that Cauchy–Schwarz inequality applied to vectors $\langle \sqrt{\alpha'_i}, \sqrt{\beta'_i} \rangle$ and $\langle \sqrt{\beta'_p}, \sqrt{\alpha'_p} \rangle$ in \mathbb{R}^2 gives

$$\sqrt{\alpha'_i \beta'_p} + \sqrt{\alpha'_p \beta'_i} \leq \sqrt{\omega'_i \omega'_p},$$

which, together with (4.7)–(4.8) leads to

$$\omega_j \leq \frac{1}{2} \sqrt{(\omega'_j)^2 - (\theta'_j)^2} + \sum_{l=0}^{\infty} \sqrt{\omega'_{j+l+1} \omega'_{j-l-1}} \quad (4.11)$$

and

$$\frac{1}{2}(\omega_j + \theta_j) \leq \lambda_0^2 \left(\sum_{l=0}^{\infty} \sqrt{\omega'_{j+l+1} \omega'_{j-l-1}} \right), \quad (4.12)$$

where λ_0 is as in (4.1) and ω'_i, θ'_i denote the right-hand derivatives.

The functions ω_j and θ_j satisfy the following properties: (i) ω_j and θ_j are absolutely continuous on $[0, \infty)$, and the right-hand derivatives ω'_j and θ'_j exist everywhere; (ii) ω_j and ω'_j are non-negative and non-increasing; (iii) ω_j is convex; (iv) ω'_j is bounded for all $j \geq k$; (v) $\omega_j(0) = \theta_j(0) = 0$; and (vi) $|\theta_j(T)| \leq \omega_j(T)$ and $|\theta'_j(T)| \leq \omega'_j(T)$ for all $T \in [0, \infty)$.

In Section IV.3 of [12] it was shown that if ω_j and θ_j are functions satisfying the above described properties (i)–(vi) and the system (4.11)–(4.12), then $\omega_j \equiv 0$ for all $j = 1, 2, \dots, k$. In particular, we have $\omega_k(T) = 0$, for all $T \in [0, \infty)$, and hence $f_k = 0$. Going back to (4.4), we get $u = 0$, and this concludes the proof of essential self-adjointness of H^k on $C_c^\infty(E)$. The essential self-adjointness of H^2, H^3, \dots , and H^{k-1} on $C_c^\infty(E)$ follows by Proposition 4.2. \square

5. PROOF OF THEOREM 2

We adapt the proof of Theorem 1.1 in [13] to our type of operator. By assumption (2.6) it follows that

$$((\Delta_{M,\mu} + q - C + 1)u, u) \geq \|u\|^2, \quad \text{for all } u \in C_c^\infty(M). \quad (5.1)$$

Since (5.1) is satisfied and since M is non-compact and g^{TM} is geodesically complete, a result of Agmon [1] (see also Proposition III.6.2 in [12]) guarantees the existence of a function $\gamma \in C^\infty(M)$ such that $\gamma(x) > 0$ for all $x \in M$, and

$$(\Delta_{M,\mu} + q - C + 1)\gamma = \gamma. \quad (5.2)$$

We now use the function γ to transform the operator $H = \nabla^* \nabla + V$. Let $L_{\mu_1}^2(E)$ be the space of square integrable sections of E with inner product $(\cdot, \cdot)_{\mu_1}$ as in (2.1), where $d\mu$ is replaced by $d\mu_1 := \gamma^2 d\mu$. For clarity, we denote $L^2(E)$ from Section 2.1 by $L_\mu^2(E)$. In what follows, the formal adjoints of ∇ with respect to inner products $(\cdot, \cdot)_\mu$ and $(\cdot, \cdot)_{\mu_1}$ will be denoted by $\nabla^{*,\mu}$ and ∇^{*,μ_1} , respectively. It is easy to check that the map $T_\gamma: L_\mu^2(E) \rightarrow L_{\mu_1}^2(E)$ defined by $Tu := \gamma^{-1}u$ is unitary. Furthermore, under the change of variables $u \mapsto \gamma^{-1}u$, the differential expression $H = \nabla^{*,\mu} \nabla + V$ gets transformed into $H_1 := \gamma^{-1}H\gamma$. Since T is unitary, the essential self-adjointness of $H^k|_{C_c^\infty(E)}$ in $L_\mu^2(E)$ is equivalent to essential self-adjointness of $(H_1)^k|_{C_c^\infty(E)}$ in $L_{\mu_1}^2(E)$.

In the sequel, we will show that H_1 has the following form:

$$H_1 = \nabla^{*,\mu_1} \nabla + \tilde{V}, \quad (5.3)$$

with

$$\tilde{V}(x) := \frac{\Delta_{M,\mu}\gamma}{\gamma} \text{Id}(x) + V(x).$$

To see this, let $w, z \in C_c^\infty(E)$ and consider

$$\begin{aligned} (H_1 w, z)_{\mu_1} &= \int_M \langle \gamma^{-1} H(\gamma w), z \rangle \gamma^2 d\mu = \int_M \langle H(\gamma w), \gamma z \rangle d\mu = (H(\gamma w), \gamma z)_\mu \\ &= (\nabla(\gamma w), \nabla(\gamma z))_\mu + (V\gamma w, \gamma z)_\mu = (\gamma^2 \nabla w, \nabla z)_\mu + (d\gamma \otimes w, d\gamma \otimes z)_{L_\mu^2(T^*M \otimes E)} \\ &\quad + (\gamma \nabla w, d\gamma \otimes z)_{L_\mu^2(T^*M \otimes E)} + (d\gamma \otimes w, \gamma \nabla z)_{L_\mu^2(T^*M \otimes E)} + (V\gamma w, \gamma z)_\mu. \end{aligned} \quad (5.4)$$

Setting $\xi := d(\gamma^2/2) \in T^*M$ and using equation (1.34) in Appendix C of [33] we have

$$(\gamma \nabla w, d\gamma \otimes z)_{L_\mu^2(T^*M \otimes E)} = (\nabla w, \xi \otimes z)_{L_\mu^2(T^*M \otimes E)} = (\nabla_X w, z)_\mu, \quad (5.5)$$

where X is the vector field associated with $\xi \in T^*M$ via the metric g^{TM} .

Furthermore, by equation (1.35) in Appendix C of [33] we have

$$\begin{aligned} (d\gamma \otimes w, \gamma \nabla z)_{L_\mu^2(T^*M \otimes E)} &= (\xi \otimes w, \nabla z)_{L_\mu^2(T^*M \otimes E)} = (\nabla^{*,\mu}(\xi \otimes w), z)_\mu \\ &= -(\text{div}_\mu(X)w, z)_\mu - (\nabla_X w, z)_\mu, \end{aligned} \quad (5.6)$$

where, in local coordinates x^1, x^2, \dots, x^n , for $X = X^j \frac{\partial}{\partial x^j}$, with Einstein summation convention,

$$\text{div}_\mu(X) := \frac{1}{\kappa} \left(\frac{\partial}{\partial x^j} (\kappa X^j) \right).$$

(Recall that $d\mu = \kappa(x) dx^1 dx^2 \dots dx^n$, where $\kappa(x)$ is a positive C^∞ -density.) Since $X^j = (g^{TM})^{jl} \left(\gamma \frac{\partial \gamma}{\partial x^j} \right)$, we have

$$\operatorname{div}_\mu(X) = |d\gamma|^2 - \gamma(\Delta_{M,\mu}\gamma), \quad (5.7)$$

where $|d\gamma(x)|$ is the norm of $d\gamma(x) \in T_x^*M$ induced by g^{TM} , and $\Delta_{M,\mu}$ is as in (1.1) with metric g^{TM} . Combining (5.4)–(5.7) and noting that

$$(d\gamma \otimes w, d\gamma \otimes z)_{L_\mu^2(T^*M \otimes E)} = \int_M |d\gamma|^2 \langle w, z \rangle d\mu,$$

we obtain

$$\begin{aligned} (H_1 w, z)_{\mu_1} &= \int_M \langle \nabla w, \nabla z \rangle \gamma^2 d\mu + \int_M \langle Vw, z \rangle \gamma^2 d\mu + \int_M \gamma(\Delta_{M,\mu}\gamma) \langle w, z \rangle d\mu \\ &= (\nabla w, \nabla z)_{L_{\mu_1}^2(T^*M \otimes E)} + (Vw, z)_{\mu_1} + (\gamma^{-1}(\Delta_{M,\mu}\gamma)w, z)_{\mu_1} \\ &= (\nabla^{*,\mu_1} \nabla w, z)_{\mu_1} + (Vw, z)_{\mu_1} + (\gamma^{-1}(\Delta_{M,\mu}\gamma)w, z)_{\mu_1}, \end{aligned} \quad (5.8)$$

which shows (5.3).

By (2.5) and (5.2) it follows that

$$\tilde{V}(x) = \frac{\Delta_{M,\mu}\gamma}{\gamma} \operatorname{Id}(x) + V(x) \geq (C-1)\operatorname{Id}(x), \quad \text{for all } x \in M,$$

where C is as in (2.6). Thus, by Theorem 1 the operator $(H_1)^k|_{C_c^\infty(E)}$ is essentially self-adjoint in $L_{\mu_1}^2(E)$ for all $k \in \mathbb{Z}_+$. \square

6. PROOF OF THEOREM 3

Throughout the section, we assume that the hypotheses of Theorem 3 are satisfied. In subsequent discussion, the notation \widehat{D} is as in (3.1) and the operators H_{\min} and H_{\max} are as in Section 4.1. We begin with the following lemma, whose proof is a direct consequence of the definition of H_{\max} and local elliptic regularity.

Lemma 6.1. *Under the assumption $V \in L_{\operatorname{loc}}^\infty(\operatorname{End} E)$, we have the following inclusion: $\operatorname{Dom}(H_{\max}) \subset W_{\operatorname{loc}}^{2,2}(E)$.*

The proof of the next lemma is given in Lemma 8.10 of [5].

Lemma 6.2. *For any $u \in \operatorname{Dom}(H_{\max})$ and any Lipschitz function with compact support $\psi: M \rightarrow \mathbb{R}$, we have:*

$$(D(\psi u), D(\psi u)) + (V\psi u, \psi u) = \operatorname{Re}(\psi H u, \psi u) + \|\widehat{D}(d\psi)u\|^2. \quad (6.1)$$

Corollary 6.3. *Let H be as in (2.3), let $u \in L^2(E)$ be a weak solution of $Hu = 0$, and let $\psi: M \rightarrow \mathbb{R}$ be a Lipschitz function with compact support. Then*

$$(\psi u, H(\psi u)) = \|\widehat{D}(d\psi)u\|^2, \quad (6.2)$$

where (\cdot, \cdot) on the left-hand side denotes the duality between $W_{\operatorname{loc}}^{1,2}(E)$ and $W_{\operatorname{comp}}^{-1,2}(E)$.

Proof. Since $u \in L^2(E)$ and $Hu = 0$, we have $u \in \text{Dom}(H_{\max}) \subset W_{\text{loc}}^{2,2}(E) \subset W_{\text{loc}}^{1,2}(E)$, where the first inclusion follows by Lemma 6.1. Since ψ is a Lipschitz compactly supported function, we get $\psi u \in W_{\text{comp}}^{1,2}(E)$ and, hence, $H(\psi u) \in W_{\text{comp}}^{-1,2}(E)$. Now the equality (6.2) follows from (6.1), the assumption $Hu = 0$, and

$$(\psi u, H(\psi u)) = (\psi u, D^* D(\psi u)) + (V\psi u, \psi u) = (D(\psi u), D(\psi u)) + (V\psi u, \psi u),$$

where in the second equality we used integration by parts; see Lemma 8.8 in [5]. Here, the two leftmost symbols (\cdot, \cdot) denote the duality between $W_{\text{comp}}^{1,2}(E)$ and $W_{\text{loc}}^{-1,2}(E)$, while the remaining ones stand for L^2 -inner products. \square

The key ingredient in the proof of Theorem 3 is the Agmon-type estimate given in the next lemma, whose proof, inspired by an idea of [25], is based on the technique developed in [10] for magnetic Laplacians on an open set with compact boundary in \mathbb{R}^n .

Lemma 6.4. *Let $\lambda \in \mathbb{R}$ and let $v \in L^2(E)$ be a weak solution of $(H - \lambda)v = 0$. Assume that there exists a constant $c_1 > 0$ such that, for all $u \in W_{\text{comp}}^{1,2}(E)$,*

$$(u, (H - \lambda)u) \geq \lambda_0^2 \int_M \max\left(\frac{1}{r(x)^2}, 1\right) |u(x)|^2 d\mu(x) + c_1 \|u\|^2, \quad (6.3)$$

where $r(x)$ is as in (2.7), λ_0 is as in (2.2), the symbol (\cdot, \cdot) on the left-hand side denotes the duality between $W_{\text{comp}}^{1,2}(E)$ and $W_{\text{loc}}^{-1,2}(E)$, and $|\cdot|$ is the norm in the fiber E_x .

Then, the following equality holds: $v = 0$.

Proof. Let ρ and R be numbers satisfying $0 < \rho < 1/2$ and $1 < R < +\infty$. For any $\varepsilon > 0$, we define the function $f_\varepsilon: M \rightarrow \mathbb{R}$ by $f_\varepsilon(x) = F_\varepsilon(r(x))$, where $r(x)$ is as in (2.7) and $F_\varepsilon: [0, \infty) \rightarrow \mathbb{R}$ is the continuous piecewise affine function defined by

$$F_\varepsilon(s) = \begin{cases} 0 & \text{for } s \leq \varepsilon \\ \rho(s - \varepsilon)/(\rho - \varepsilon) & \text{for } \varepsilon \leq s \leq \rho \\ s & \text{for } \rho \leq s \leq 1 \\ 1 & \text{for } 1 \leq s \leq R \\ R + 1 - s & \text{for } R \leq s \leq R + 1 \\ 0 & \text{for } s \geq R + 1. \end{cases}$$

Let us fix $x_0 \in M$. For any $\alpha > 0$, we define the function $p_\alpha: M \rightarrow \mathbb{R}$ by

$$p_\alpha(x) = P_\alpha(d_{g_{TM}}(x_0, x)),$$

where $P_\alpha: [0, \infty) \rightarrow \mathbb{R}$ is the continuous piecewise affine function defined by

$$P_\alpha(s) = \begin{cases} 1 & \text{for } s \leq 1/\alpha \\ -\alpha s + 2 & \text{for } 1/\alpha \leq s \leq 2/\alpha \\ 0 & \text{for } s \geq 2/\alpha. \end{cases}$$

Since $\widehat{d}_{g_{TM}}(x_0, x) \leq d_{g_{TM}}(x_0, x)$, it follows that the support of $f_\varepsilon p_\alpha$ is contained in the set $B_\alpha := \{x \in M: \widehat{d}_{g_{TM}}(x_0, x) \leq 2/\alpha\}$. By assumption (A1) we know that \widehat{M} is a geodesically complete Riemannian manifold. Hence, by Hopf–Rinow Theorem the set B_α is compact. Therefore, the

support of $f_\varepsilon p_\alpha$ is compact. Additionally, note that $f_\varepsilon p_\alpha$ is a β -Lipschitz function (with respect to the distance corresponding to the metric g^{TM}) with $\beta = \frac{\rho}{\rho - \varepsilon} + \alpha$.

Since $v \in L^2(E)$ and $(H - \lambda)v = 0$, we have $v \in \text{Dom}(H_{\max}) \subset W_{\text{loc}}^{2,2}(E) \subset W_{\text{loc}}^{1,2}(E)$, where the first inclusion follows by Lemma 6.1. Since $f_\varepsilon p_\alpha$ is a Lipschitz compactly supported function, we get $f_\varepsilon p_\alpha v \in W_{\text{comp}}^{1,2}(E)$ and, hence, $((H - \lambda)(f_\varepsilon p_\alpha v)) \in W_{\text{comp}}^{-1,2}(E)$.

Using (2.2) we have

$$\|\widehat{D}(d(f_\varepsilon p_\alpha))v\|^2 \leq \lambda_0^2 \int_M |d(f_\varepsilon p_\alpha)(x)|^2 |v(x)|^2 d\mu(x), \quad (6.4)$$

where $|d(f_\varepsilon p_\alpha)(x)|$ is the norm of $d(f_\varepsilon p_\alpha)(x) \in T_x^*M$ induced by g^{TM} .

By Corollary 6.3 with $H - \lambda$ in place of H and the inequality (6.4), we get

$$(f_\varepsilon p_\alpha v, (H - \lambda)(f_\varepsilon p_\alpha v)) \leq \lambda_0^2 \left(\frac{\rho}{\rho - \varepsilon} + \alpha \right)^2 \|v\|^2. \quad (6.5)$$

On the other hand, using the definitions of f_ε and p_α and the assumption (6.3) we have

$$(f_\varepsilon p_\alpha v, (H - \lambda)(f_\varepsilon p_\alpha v)) \geq \lambda_0^2 \int_{S_{\rho,R,\alpha}} |v(x)|^2 d\mu(x) + c_1 \|f_\varepsilon p_\alpha v\|^2, \quad (6.6)$$

where

$$S_{\rho,R,\alpha} := \{x \in M : \rho \leq r(x) \leq R \text{ and } d_{g^{TM}}(x_0, x) \leq 1/\alpha\}.$$

In (6.6) and (6.5), the symbol (\cdot, \cdot) stands for the duality between $W_{\text{comp}}^{1,2}(E)$ and $W_{\text{loc}}^{-1,2}(E)$. We now combine (6.6) and (6.5) to get

$$\lambda_0^2 \int_{S_{\rho,R,\alpha}} |v(x)|^2 d\mu(x) + c_1 \|f_\varepsilon p_\alpha v\|^2 \leq \lambda_0^2 \left(\frac{\rho}{\rho - \varepsilon} + \alpha \right)^2 \|v\|^2.$$

We fix ρ , R , and ε , and let $\alpha \rightarrow 0+$. After that we let $\varepsilon \rightarrow 0+$. The last step is to do $\rho \rightarrow 0+$ and $R \rightarrow +\infty$. As a result, we get $v = 0$. \square

End of the proof of Theorem 3. Using integration by parts (see Lemma 8.8 in [5]), we have

$$(u, Hu) = (u, D^*Du) + (Vu, u) = (Du, Du) + (Vu, u) \geq (Vu, u), \quad \text{for all } u \in W_{\text{comp}}^{1,2}(E),$$

where the two leftmost symbols (\cdot, \cdot) denote the duality between $W_{\text{comp}}^{1,2}(E)$ and $W_{\text{loc}}^{-1,2}(E)$, while the remaining ones stand for L^2 -inner products. Hence, by assumption (2.8) we get:

$$\begin{aligned} (u, (H - \lambda)u) &\geq \lambda_0^2 \int_M \frac{1}{r(x)^2} |u(x)|^2 d\mu(x) - (\lambda + C) \|u\|^2 \\ &\geq \lambda_0^2 \int_M \max\left(\frac{1}{r(x)^2}, 1\right) |u(x)|^2 d\mu(x) - (\lambda + C + 1) \|u\|^2. \end{aligned} \quad (6.7)$$

Choosing, for instance, $\lambda = -C - 2$ in (6.7) we get the inequality (6.3) with $c_1 = 1$.

Thus, $H_{\min} - \lambda$ with $\lambda = -C - 2$ is a symmetric operator satisfying $(u, (H_{\min} - \lambda)u) \geq \|u\|^2$, for all $u \in C_c^\infty(E)$. In this case, it is known (see Theorem X.26 in [29]) that the essential self-adjointness of $H_{\min} - \lambda$ is equivalent to the following statement: if $v \in L^2(E)$ satisfies

$(H - \lambda)v = 0$, then $v = 0$. Thus, by Lemma 6.4, the operator $(H_{\min} - \lambda)$ is essentially self-adjoint. Hence, H_{\min} is essentially self-adjoint. \square

REFERENCES

- [1] Agmon, S.: On positivity and decay of solutions of second order elliptic equations on Riemannian manifolds. In: *Methods of functional analysis and theory of elliptic equations* (Naples, 1982) Liguori, Naples (1983) pp. 19–52
- [2] Bandara, L.: Density problems on vector bundles and manifolds. *Proc. Amer. Math. Soc.* **142** (2014) 2683–2695
- [3] Braverman, M.: On self-adjointness of Schrödinger operator on differential forms. *Proc. Amer. Math. Soc.* **126** (1998) 617–623
- [4] Braverman, M., Cecchini, S.: Spectral theory of von Neumann algebra valued differential operators over non-compact manifolds. *arXiv:1503.02998*
- [5] Braverman, M., Milatovic, O., Shubin, M.: Essential self-adjointness of Schrödinger-type operators on manifolds. *Russian Math. Surveys* **57** (2002) 641–692
- [6] Brusentsev, A. G.: Self-adjointness of elliptic differential operators in $L_2(G)$ and correcting potentials. *Trans. Moscow Math. Soc.* (2004) 31–61
- [7] Chernoff, P.: Essential self-adjointness of powers of generators of hyperbolic equations. *J. Funct. Analysis* **12** (1973) 401–414
- [8] Chernoff, P.: Schrödinger and Dirac operators with singular potentials and hyperbolic equations. *Pacific J. Math* **72** (1977) 361–382
- [9] Chumak A. A.: Self-adjointness of the Beltrami-Laplace operator on a complete paracompact manifold without boundary. *Ukrainian Math. Journal* **25** (1973) 784–791 (Russian)
- [10] Colin de Verdière, Y., Truc, F: Confining quantum particles with a purely magnetic field. *Ann. Inst. Fourier (Grenoble)* **60** (7) (2010) 2333–2356
- [11] Cordes, H. O.: Self-adjointness of powers of elliptic operators on non-compact manifolds. *Math. Annalen* **195** (1972) 257–272
- [12] Cordes, H. O.: *Spectral Theory of Linear Differential Operators and Comparison Algebras*. London Math. Soc., Lecture Notes Series **76**, Cambridge University Press (1987)
- [13] Cordes, H. O.: On essential selfadjointness of powers and comparison algebras. *Festschrift on the occasion of the 70th birthday of Shmuel Agmon. J. Anal. Math.* **58** (1992) 61–97
- [14] Cycon, H. L., Froese, R. G., Kirsch, W., Simon, B.: *Schrödinger Operators with Applications to Quantum Mechanics and Global Geometry*. Texts and Monographs in Physics, Springer-Verlag (1987)
- [15] Gaffney, M.: A special Stokes’s theorem for complete Riemannian manifolds. *Ann. of Math.* **60** (1954) 140–145
- [16] Gaffney, M.: Hilbert space methods in the theory of harmonic integrals. *Trans. Amer. Math. Soc.* **78** (1955) 426–444
- [17] Grigor’yan, A., Masamune, J.: Parabolicity and stochastic completeness of manifolds in terms of Green formula. *J. Math. Pures Appl.* (9) **100** (2013) 607–632.
- [18] Grummt, R., Kolb, M.: Essential selfadjointness of singular magnetic Schrödinger operators on Riemannian manifolds. *J. Math. Anal. Appl.* **388** (2012) 480–489
- [19] Güneysu, B., Post, O.: Path integrals and the essential self-adjointness of differential operators on noncompact manifolds. *Math. Z.* **275** (2013) 331–348
- [20] Kato, T.: *Perturbation Theory for Linear Operators*. Springer-Verlag, Berlin (1980)
- [21] Lesch, M.: Essential self-adjointness of symmetric linear relations associated to first order systems. *Journées Équations aux Dérivées Partielles (La Chapelle sur Erdre) Univ. Nantes, Exp. No. X* (2000) 18pp
- [22] Masamune, J.: Essential self-adjointness of Laplacians on Riemannian manifolds with fractal boundary. *Comm. Partial Differential Equations* **24** (1999) 749–757

- [23] Masamune, J.: Analysis of the Laplacian of an incomplete manifold with almost polar boundary. *Rend. Mat. Appl.* (7) **25** (2005) 109–126
- [24] Masamune, J.: Conservative principle for differential forms. *Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei* (9) Mat. Appl. **18** (2007) 351–358
- [25] Nenciu, G., Nenciu, I.: On confining potentials and essential self-adjointness for Schrödinger operators on bounded domains in \mathbb{R}^n . *Ann. Henri Poincaré* **10** (2009) 377–394
- [26] Nenciu, G., Nenciu, I.: On essential self-adjointness for magnetic Schrödinger and Pauli operators on the unit disc in \mathbb{R}^2 . *Lett. Math. Phys.* **98** (2011) 207–223
- [27] Oleinik, I.: On the essential self-adjointness of the Schrödinger operator on complete Riemannian manifolds. *Math. Notes* **54** (1993) 934–939
- [28] Oleinik, I.: On a connection between classical and quantum-mechanical completeness of the potential at infinity on a complete Riemannian manifold. *Math. Notes* **55** (1994) 380–386
- [29] Reed, M., Simon, B.: *Methods of Modern Mathematical Physics II: Fourier Analysis, Self-Adjointness*. Academic Press, New York (1975)
- [30] Roelcke, W.: Über den Laplace-Operator auf Riemannschen Mannigfaltigkeiten mit diskontinuierlichen Gruppen. *Math. Nachr.* **21** (1960) 131–149 (German)
- [31] Shubin, M. A.: Spectral theory of elliptic operators on noncompact manifolds. *Astérisque* **207** (1992) 35–108
- [32] Shubin, M.: Essential self-adjointness for magnetic Schrödinger operators on non-compact manifolds. In: *Séminaire Équations aux Dérivées Partielles (Polytechnique) (1998-1999)*, Exp. No. XV, Palaiseau (1999) pp. XV-1–XV-22
- [33] Taylor, M.: *Partial Differential Equations II: Qualitative Studies of Linear Equations*, Springer, New York (1996)

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF NORTH FLORIDA, JACKSONVILLE, FL 32224, USA.

E-mail address: `omilatov@unf.edu`

GRENOBLE UNIVERSITY, INSTITUT FOURIER, UNITÉ MIXTE DE RECHERCHE CNRS-UJF 5582, BP 74, 38402-SAINT MARTIN D'HÈRES CEDEX, FRANCE.

E-mail address: `francoise.truc@ujf-grenoble.fr`